

# ORIGINS

**E.P. Wigner**, Conference on Neutron Physics by Time of Flight, November **1956**

**P.W. Anderson**, “*Absence of Diffusion in Certain Random Lattices*”; Phys.Rev., **1958**, v.109, p.1492

**L.D. Landau**, “*Fermi-Liquid Theory*” Zh. Exp. Teor. Fiz., **1956**, v.30, p.1058

**J. Bardeen, L.N. Cooper & J. Schriffer**, “*Theory of Superconductivity*”; Phys.Rev., **1957**, v.108, p.1175.

What does it mean - **non-Fermi liquid** ?


Q : What is the difference between **Fermi-liquid** and **non-Fermi liquid** ?

A : The difference is the same as between **bananas** and **non-bananas**.

What does it mean **Fermi liquid** ?

# *Fermi Liquid*

- *Fermi statistics*
- *Low temperatures*
- *Not too strong interactions*
- *Translation invariance*



*Fermi  
Liquid*

*What does it mean?*

- *Fermi statistics*
- *Low temperatures*
- *Not too strong interactions*
- *Translation invariance*



*Fermi  
Liquid*

*It means that*

1. *Excitations are similar to the excitations in a Fermi-gas:*
  - a) *the same quantum numbers – momentum, spin  $\frac{1}{2}$ , charge  $e$*
  - b) *decay rate is small as compared with the excitation energy*
2. *Substantial renormalizations. For example, in a Fermi gas*

$$\partial n / \partial \mu, \quad \gamma = c / T, \quad \chi / g \mu_B$$

*are all equal to the one-particle density of states  $\nu$ .  
These quantities are different in a Fermi liquid*

## Signatures of the Fermi - Liquid state ?!

### 1. Resistivity is proportional to $T^2$ :

L.D. Landau & I.Ya. Pomeranchuk “*To the properties of metals at very low temperatures*”; Zh.Exp.Teor.Fiz., **1936**, v.10, p.649

...The increase of the resistance caused by the interaction between the electrons is proportional to  $T^2$  and at low temperatures exceeds the **usual** resistance, which is proportional to  $T^5$ .

... the sum of the momenta of the interaction electrons **can change** by an **integer number of the periods of the reciprocal lattice**. Therefore the momentum increase caused by the electric field can be destroyed by the interaction between the electrons, not only by the thermal oscillations of the lattice.

## *Signatures of the Fermi - Liquid state ?!*

1. *Resistivity is proportional to  $T^2$  :*

**L.D. Landau & I.Ya. Pomeranchuk** “*To the properties of metals at very low temperatures*”; **Zh.Exp.Teor.Fiz.**, **1936**, v.10, p.649

*Umklapp* electron – electron scattering dominates the charge transport (?!)


# Signatures of the Fermi - Liquid state ?!

1. Resistivity is proportional to  $T^2$  :

L.D. Landau & I.Ya. Pomeranchuk “To the properties of metals at very low temperatures”; Zh.Exp.Teor.Fiz., 1936, v.10, p.649

*Umklapp* electron – electron scattering dominates the charge transport (?!)

2. Jump in the momentum distribution function at  $T=0$ .




# Signatures of the Fermi - Liquid state ?!

1. Resistivity is proportional to  $T^2$  :

L.D. Landau & I.Ya. Pomeranchuk “To the properties of metals at very low temperatures”; Zh.Exp.Teor.Fiz., 1936, v.10, p.649

*Umklapp* electron – electron scattering dominates the charge transport (?!)

2. Jump in the momentum distribution function at  $T=0$ .



2a. Pole in the one-particle Green function

$$G(\varepsilon, \vec{p}) = \frac{Z}{i\varepsilon_n - \xi(\vec{p})}$$

Fermi liquid =  $0 < Z < 1$  (?!)



# Landau Fermi - Liquid theory

*Momentum*

$\vec{p}$

*Momentum distribution*

$n(\vec{p})$

*Total energy*

$E\{n(\vec{p})\}$

*Quasiparticle energy*

$\xi(\vec{p}) \equiv \delta E / \delta n(\vec{p})$

*Landau f-function*

$f(\vec{p}, \vec{p}') \equiv \delta \xi(\vec{p}) / \delta n(\vec{p}')$

# Landau Fermi - Liquid theory

*Momentum*

$$\vec{p}$$

*Momentum distribution*

$$n(\vec{p})$$

*Total energy*

$$E\{n(\vec{p})\}$$

*Quasiparticle energy*

$$\xi(\vec{p}) \equiv \delta E / \delta n(\vec{p})$$

*Landau f-function*

$$f(\vec{p}, \vec{p}') \equiv \delta \xi(\vec{p}) / \delta n(\vec{p}')$$

Q:

Can *Fermi - liquid* survive without the *momenta*

Does it make sense to speak about the *Fermi - liquid* state in the presence of a *quenched disorder*

?

Q


• Does it make sense to speak about the *Fermi - liquid* state in the presence of a *quenched disorder*

?

1. Momentum *is not* a good quantum number – the momentum uncertainty is inverse proportional to the *elastic mean free path*,  $l$ . The step in the momentum distribution function is broadened by this uncertainty


Q • Does it make sense to speak about the *Fermi – liquid state* in the presence of a *quenched disorder* ?

1. Momentum *is not* a good quantum number – the momentum uncertainty is inverse proportional to the *elastic mean free path,  $l$* . The step in the momentum distribution function is broadened by this uncertainty



Q: Does it make sense to speak about the *Fermi - liquid* state in the presence of a *quenched disorder* ?

1. Momentum *is not* a good quantum number – the momentum uncertainty is inverse proportional to the *elastic mean free path*,  $l$ . The step in the momentum distribution function is broadened by this uncertainty




2. Neither resistivity nor its temperature dependence is determined by the *umklapp processes* and thus does not behave as  $T^2$
3. Sometimes (e.g., for random quenched magnetic field) the disorder averaged one-particle Green function even without interactions *does not have a pole* as a function of the energy,  $\epsilon$ . The residue,  $Z$ , makes no sense.

*Nevertheless even in the presence of the disorder*

- I. Excitations are *similar* to the excitations in a disordered *Fermi-gas*.
- II. Small decay rate
- III. Substantial renormalizations

# Quantum Dot



1. Disorder ( $\times$  impurities)
  2. Complex geometry
  3.  $e-e$  interactions
- } *chaotic one-particle motion*

## Realizations:


- Metallic clusters
- Gate determined confinement in 2D gases (e.g. *GaAs/AlGaAs*)
- Carbon nanotubes
- 
-

# One-particle problem (*Thouless, 1972*)


Energy scales

## 1. Mean level spacing

$$\delta_1 = 1/\nu \times L^d$$



energy



$L$  is the system size;

$d$  is the number of dimensions

## 2. Thouless energy

$$E_T = hD/L^2$$

$D$  is the diffusion const

$E_T$  has a meaning of the *inverse diffusion time* of the traveling through the system or the *escape rate* (for open systems)

$$g = E_T / \delta_1$$

dimensionless  
*Thouless*  
conductance

$$g = Gh/e^2$$

# Zero Dimensional Fermi Liquid

Finite  
System



Thouless  
energy  $E_T$

$$\varepsilon \ll E_T \xrightarrow{\text{def}} 0D$$

At the same time, we want the typical energies,  $\varepsilon$ , to exceed the mean level spacing,  $\delta_1$ :

$$\delta_1 \ll \varepsilon \ll E_T$$

$$g \equiv \frac{E_T}{\delta_1} \gg 1$$



# Thouless Conductance and One-particle Quantum Mechanics



*Localized states*  
*Insulator*


*Extended states*  
*Metal*

**Poisson spectral statistics**


**Wigner-Dyson spectral statistics**

$N \times N$   
*Random Matrices*

*Quantum Dots with dimensionless conductance  $g$*



*The same statistics of the random spectra and one-particle wave functions (eigenvectors)*



# Two-Body Interactions

$$|\alpha, \sigma\rangle$$

*Set of one particle states.  $\sigma$  and  $\alpha$  label correspondingly *spin* and *orbit*.*

# Two-Body Interactions

$|\alpha, \sigma\rangle$

Set of one particle states.  $\sigma$  and  $\alpha$  label correspondingly *spin* and *orbit*.

$$\hat{H}_0 = \sum_{\alpha} \varepsilon_{\alpha} a_{\alpha, \sigma}^{\dagger} a_{\alpha, \sigma} \quad \hat{H}_{\text{int}} = \sum_{\substack{\alpha, \beta, \gamma, \delta \\ \sigma, \sigma'}} M_{\alpha\beta\gamma\delta} a_{\alpha, \sigma}^{\dagger} a_{\beta, \sigma'}^{\dagger} a_{\gamma, \sigma} a_{\delta, \sigma'}$$

$\varepsilon_{\alpha}$  -one-particle orbital energies

$M_{\alpha\beta\gamma\delta}$  -interaction matrix elements

# Two-Body Interactions


$|\alpha, \sigma\rangle$

Set of one particle states.  $\sigma$  and  $\alpha$  label correspondingly *spin* and *orbit*.

$$\hat{H}_0 = \sum_{\alpha} \epsilon_{\alpha} a_{\alpha, \sigma}^{\dagger} a_{\alpha, \sigma} \quad \hat{H}_{\text{int}} = \sum_{\substack{\alpha, \beta, \gamma, \delta \\ \sigma, \sigma'}} M_{\alpha\beta\gamma\delta} a_{\alpha, \sigma}^{\dagger} a_{\beta, \sigma'}^{\dagger} a_{\gamma, \sigma} a_{\delta, \sigma'}$$

$\epsilon_{\alpha}$  -one-particle orbital energies

$M_{\alpha\beta\gamma\delta}$  -interaction matrix elements



# Two-Body Interactions

$|\alpha, \sigma\rangle$

Set of one particle states.  $\sigma$  and  $\alpha$  label correspondingly *spin* and *orbit*.

$$\hat{H}_0 = \sum_{\alpha} \varepsilon_{\alpha} a_{\alpha, \sigma}^{\dagger} a_{\alpha, \sigma} \quad \hat{H}_{\text{int}} = \sum_{\substack{\alpha, \beta, \gamma, \delta \\ \sigma, \sigma'}} M_{\alpha\beta\gamma\delta} a_{\alpha, \sigma}^{\dagger} a_{\beta, \sigma'}^{\dagger} a_{\gamma, \sigma} a_{\delta, \sigma'}$$

$\varepsilon_{\alpha}$  -one-particle orbital energies

$M_{\alpha\beta\gamma\delta}$  -interaction matrix elements

# Two-Body Interactions

$|\alpha, \sigma\rangle$

Set of one particle states.  $\sigma$  and  $\alpha$  label correspondingly *spin* and *orbit*.

$$\hat{H}_0 = \sum_{\alpha} \varepsilon_{\alpha} a_{\alpha, \sigma}^{\dagger} a_{\alpha, \sigma} \quad \hat{H}_{\text{int}} = \sum_{\substack{\alpha, \beta, \gamma, \delta \\ \sigma, \sigma'}} M_{\alpha\beta\gamma\delta} a_{\alpha, \sigma}^{\dagger} a_{\beta, \sigma'}^{\dagger} a_{\gamma, \sigma} a_{\delta, \sigma'}$$

$\varepsilon_{\alpha}$  -one-particle orbital energies

$M_{\alpha\beta\gamma\delta}$  -interaction matrix elements

*Nuclear  
Physics*

$\varepsilon_{\alpha}$

are taken from the *shell model*

$M_{\alpha\beta\gamma\delta}$

are assumed to be *random*

# Two-Body Interactions

$$|\alpha, \sigma\rangle$$

Set of one particle states.  $\sigma$  and  $\alpha$  label correspondingly *spin* and *orbit*.

$$\hat{H}_0 = \sum_{\alpha} \varepsilon_{\alpha} a_{\alpha, \sigma}^{\dagger} a_{\alpha, \sigma} \quad \hat{H}_{\text{int}} = \sum_{\substack{\alpha, \beta, \gamma, \delta \\ \sigma, \sigma'}} M_{\alpha\beta\gamma\delta} a_{\alpha, \sigma}^{\dagger} a_{\beta, \sigma'}^{\dagger} a_{\gamma, \sigma} a_{\delta, \sigma'}$$

$\varepsilon_{\alpha}$  -one-particle orbital energies

$M_{\alpha\beta\gamma\delta}$  -interaction matrix elements

Nuclear  
Physics

$$\varepsilon_{\alpha}$$

are taken from the *shell model*

$$M_{\alpha\beta\gamma\delta}$$

are assumed to be *random*

Quantum  
Dots

$$\varepsilon_{\alpha}$$

*RANDOM*; Wigner-Dyson statistics

$$M_{\alpha\beta\gamma\delta}$$

????????

# Matrix Elements

$$\hat{H}_{\text{int}} = \sum_{\substack{\alpha, \beta, \gamma, \delta \\ \sigma, \sigma'}} M_{\alpha\beta\gamma\delta} a_{\alpha, \sigma}^+ a_{\beta, \sigma'}^+ a_{\gamma, \sigma} a_{\delta, \sigma'}$$

$$M_{\alpha\beta\gamma\delta}$$

*Diagonal* -  $\alpha, \beta, \gamma, \delta$  are equal *pairwise*

$\alpha = \gamma$  and  $\beta = \delta$  or  $\alpha = \delta$  and  $\beta = \gamma$  or  $\alpha = \beta$  and  $\gamma = \delta$

*Offdiagonal* - *otherwise*



# Matrix Elements

$$\hat{H}_{\text{int}} = \sum_{\substack{\alpha, \beta, \gamma, \delta \\ \sigma, \sigma'}} M_{\alpha\beta\gamma\delta} a_{\alpha, \sigma}^+ a_{\beta, \sigma'}^+ a_{\gamma, \sigma} a_{\delta, \sigma'}$$

$$M_{\alpha\beta\gamma\delta}$$

*Diagonal* -  $\alpha, \beta, \gamma, \delta$  are equal *pairwise*

$\alpha=\gamma$  and  $\beta=\delta$  or  $\alpha=\delta$  and  $\beta=\gamma$  or  $\alpha=\beta$  and  $\gamma=\delta$

*Offdiagonal* - *otherwise*

It turns  
out that

in the limit  $g \rightarrow \infty$

- *Diagonal* matrix elements are *much bigger* than the *offdiagonal* ones

$$M_{\text{diagonal}} \gg M_{\text{offdiagonal}}$$

- *Diagonal* matrix elements in a particular sample do not fluctuate - *selfaveraging*

# Toy model:

Short range **e-e** interactions

$$U(\vec{r}) = \frac{\lambda}{\nu} \delta(\vec{r})$$

$\lambda$  is dimensionless coupling constant  
 $\nu$  is the electron density of states

$$M_{\alpha\beta\gamma\delta} = \frac{\lambda}{\nu} \int d\vec{r} \psi_{\alpha}^*(\vec{r}) \psi_{\beta}^*(\vec{r}) \psi_{\gamma}(\vec{r}) \psi_{\delta}(\vec{r})$$

$$\psi_{\alpha}(\vec{r})$$

one-particle  
eigenfunctions

# Toy model:


Short range **e-e** interactions

$$U(\vec{r}) = \frac{\lambda}{\nu} \delta(\vec{r})$$

$\lambda$  is dimensionless coupling constant  
 $\nu$  is the electron density of states

$$M_{\alpha\beta\gamma\delta} = \frac{\lambda}{\nu} \int d\vec{r} \psi_{\alpha}^*(\vec{r}) \psi_{\beta}^*(\vec{r}) \psi_{\gamma}(\vec{r}) \psi_{\delta}(\vec{r})$$

$\psi_{\alpha}(\vec{r})$   
one-particle  
eigenfunctions



# Toy model:

Short range  $e-e$   
interactions


$$U(\vec{r}) = \frac{\lambda}{\nu} \delta(\vec{r})$$

$\lambda$  is dimensionless coupling constant;  $\nu$  is the electron density of states

$$M_{\alpha\beta\gamma\delta} = \frac{\lambda}{\nu} \int d\vec{r} \psi_{\alpha}^*(\vec{r}) \psi_{\beta}^*(\vec{r}) \psi_{\gamma}(\vec{r}) \psi_{\delta}(\vec{r})$$

$$\psi_{\alpha}(\vec{r})$$

one-particle  
eigenfunctions



$\psi_{\alpha}(\vec{r})$  is a random function  
that rapidly oscillates

$$|\psi_{\alpha}(\vec{r})|^2 \geq 0$$

$\psi_{\alpha}(\vec{r})^2 \geq 0$  as long as  
 $T$ -invariance  
is preserved

## In the limit

$$g \rightarrow \infty$$

- *Diagonal matrix elements are much bigger than the offdiagonal ones*

$$M_{\text{diagonal}} \gg M_{\text{offdiagonal}}$$

- *Diagonal matrix elements in a particular sample do not fluctuate - selfaveraging*

$$M_{\alpha\beta\alpha\beta} = \frac{\lambda}{V} \int d\vec{r} |\psi_{\alpha}(\vec{r})|^2 |\psi_{\beta}(\vec{r})|^2$$

$$|\psi_{\alpha}(\vec{r})|^2 \Rightarrow \frac{1}{\text{volume}}$$

$$M_{\alpha\beta\alpha\beta} = \lambda \delta_{\alpha\beta}$$

## In the limit

$$g \rightarrow \infty$$

- *Diagonal matrix elements are much bigger than the offdiagonal ones*

$$M_{\text{diagonal}} \gg M_{\text{offdiagonal}}$$

- *Diagonal matrix elements in a particular sample do not fluctuate - selfaveraging*

$$M_{\alpha\beta\alpha\beta} = \frac{\lambda}{V} \int d\vec{r} |\psi_{\alpha}(\vec{r})|^2 |\psi_{\beta}(\vec{r})|^2$$

$$|\psi_{\alpha}(\vec{r})|^2 \Rightarrow \frac{1}{\text{volume}}$$

$$M_{\alpha\beta\alpha\beta} = \lambda \delta_{\alpha\beta}$$

**More general:** *finite range interaction potential*  $U(\vec{r})$

$$M_{\alpha\beta\alpha\beta} = \frac{\lambda}{V} \int |\psi_{\alpha}(\vec{r}_1)|^2 |\psi_{\beta}(\vec{r}_2)|^2 U(\vec{r}_1 - \vec{r}_2) d\vec{r}_1 d\vec{r}_2$$

*The same conclusion*

# Random Matrices:

$E_\alpha$  - spectrum

$\psi_\alpha(i)$  -  $i$ -th component of  $\alpha$ -th eigenvector

$$\langle \psi_\alpha^*(i) \psi_\gamma(j) \rangle = \frac{1}{N} \delta_{\alpha\gamma} \delta_{ij}$$

$$\langle \psi_\alpha(i) \psi_\gamma(j) \rangle = \frac{2-\beta}{N} \delta_{\alpha\gamma} \delta_{ij}$$

in the limit  $N \rightarrow \infty$

Components of the different eigenvectors as well as different components of the same eigenvector are not correlated

# Berry

## Conjecture:

Exact wavefunctions at energy  $\approx \mathcal{E}_F$  in chaotic systems behave as sums of plane waves with  $|\vec{k}| \approx k_F$  and random coefficients:

$$\langle \psi_\alpha^*(\vec{r}_1) \psi_\gamma(\vec{r}_2) \rangle = \frac{\delta_{\alpha\gamma}}{V} f\left(\frac{2\pi|\vec{r}_1 - \vec{r}_2|}{\lambda}\right)$$

$$\langle \psi_\alpha(\vec{r}_1) \psi_\gamma(\vec{r}_2) \rangle = (2 - \beta) \frac{\delta_{\alpha\gamma}}{V} f\left(\frac{2\pi|\vec{r}_1 - \vec{r}_2|}{\lambda}\right)$$

$$f(x) = \Gamma\left(\frac{d}{2}\right) x^{1-d/2} J_{d/2-1}(x)$$

$d$  is # of dimensions,  
 $J_\mu(x)$  is Bessel function

**Important:**



*when  $x$  increases  $f(x)$  decays quickly enough for the integral*


$$\int_0^\infty f(x) x^{d-1} dx \quad \text{to converge}$$

**Only local correlations**



# AFRICA BILLIARD - *a conformal image of a unit circle*

$$\omega(z) = R \frac{z + bz^2 + ce^{i\delta} z^3}{\sqrt{1 + 2b^2 + 3c^3}}$$




$$b = c = 0.2;$$

$$\delta = 1.5; R = 1$$

# Distribution of the matrix elements

$$M_{\alpha\beta\gamma\delta} = \int d\vec{r} \psi_{\alpha}(\vec{r})\psi_{\beta}(\vec{r})\psi_{\alpha}^*(\vec{r})\psi_{\beta}^*(\vec{r})$$




**Open boxes -**  
levels from **20 to 30**

**Closed circuits -**  
levels from **30 to 40**

$$M_{\alpha\beta\gamma\delta} = \frac{1}{\pi} \int d\vec{r} \psi_{\alpha}(\vec{r}) \psi_{\beta}(\vec{r}) \psi_{\alpha}^*(\vec{r}) \psi_{\beta}^*(\vec{r})$$

$$V_{\alpha\beta\gamma\delta} \propto \int \frac{d\vec{r}_1 d\vec{r}_2}{|\vec{r}_1 - \vec{r}_2|} \psi_{\alpha}(\vec{r}_1) \psi_{\beta}(\vec{r}_2) \psi_{\alpha}^*(\vec{r}_1) \psi_{\beta}^*(\vec{r}_2)$$

Distribution  
function of  
diagonal  
and  
offdiagonal  
matrix  
elements



**Universal** (Random Matrix) limit - Random Matrix **symmetry** of the correlation functions:

All correlation functions are invariant under arbitrary orthogonal transformation:

$$\tilde{\psi}_\mu(\vec{r}) = \sum_\nu \int d\vec{r}_1 O_\mu^\nu(\vec{r}, \vec{r}_1) \psi_\nu(\vec{r}_1)$$

$$\int d\vec{r}_1 O_\mu^\nu(\vec{r}, \vec{r}_1) O_\nu^\eta(\vec{r}_1, \vec{r}') = \delta_{\mu\eta} \delta(\vec{r} - \vec{r}')$$

There are **only** three operators, which are quadratic in the fermion operators  $a^+$ ,  $a$ , and invariant under **RM** transformations:

$$\hat{n} = \sum_{\alpha, \sigma} a_{\alpha, \sigma}^+ a_{\alpha, \sigma}$$

**total number of particles**

$$\hat{S} = \sum_{\alpha, \sigma_1, \sigma_2} a_{\alpha, \sigma_1}^+ \vec{\sigma}_{\sigma_1, \sigma_2} a_{\alpha, \sigma_2}$$

**total spin**

$$\hat{T}^+ = \sum_{\alpha} a_{\alpha, \uparrow}^+ a_{\alpha, \downarrow}^+$$

**????**

Charge conservation  
(gauge invariance)

-no  $\hat{T}$  or  $\hat{T}^+$  only  $\hat{T} \hat{T}^+$

Invariance under  
rotations in spin space

-no  $\hat{S}$  only  $\hat{S}^2$

Therefore, in a very general case

$$\hat{H}_{\text{int}} = eV\hat{n} + E_c\hat{n}^2 + J\hat{S}^2 + \lambda_{BCS}\hat{T}^+\hat{T}.$$

**Only** three coupling constants describe **all** of the effects of e-e interactions

In a very general case **only three** coupling constants describe **all** effects of electron-electron interactions:

$$\hat{H} = \sum_{\alpha} \varepsilon_{\alpha} n_{\alpha} + \hat{H}_{\text{int}}$$

$$\hat{H}_{\text{int}} = eV\hat{n} + E_c\hat{n}^2 + J\hat{S}^2 + \lambda_{\text{BCS}}\hat{T}^+\hat{T}.$$

*I.L. Kurland, I.L.Aleiner & B.A., 2000*

*See also*

*P.W.Brouwer, Y.Oreg & B.I.Halperin, 1999*

*H.Baranger & L.I.Glazman, 1999*

*H-Y Kee, I.L.Aleiner & B.A., 1998*

In a very general case **only three** coupling constants describe **all** effects of electron-electron interactions:

$$\hat{H} = \sum_{\alpha} \varepsilon_{\alpha} n_{\alpha} + \hat{H}_{\text{int}}$$

$$\hat{H}_{\text{int}} = eV\hat{n} + E_c\hat{n}^2 + J\hat{S}^2 + \lambda_{\text{BCS}}\hat{T}^+\hat{T}.$$

For a short range interaction with a coupling constant  $\lambda$

$$E_c = \frac{\lambda\delta_1}{2} \quad J = -2\lambda\delta_1 \quad \lambda_{\text{BCS}} = \lambda\delta_1(2 - \beta)$$

where  $\delta_1$  is the one-particle mean level spacing



$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{int}}$$

$$\hat{H}_0 = \sum_{\alpha} \varepsilon_{\alpha} n_{\alpha}$$

$$\hat{H}_{\text{int}} = eV\hat{n} + E_c\hat{n}^2 + J\hat{S}^2 + \lambda_{BCS}\hat{T}^+\hat{T}.$$

$E_c$  determines the charging energy  
(Coulomb blockade)

$J$  describes the spin exchange interaction

$\lambda_{BCS}$  determines effect of superconducting-like pairing

$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{int}}$$

$$\hat{H}_0 = \sum_{\alpha} \varepsilon_{\alpha} n_{\alpha}$$



$$\hat{H}_{\text{int}} = eV\hat{n} + E_c\hat{n}^2 + J\hat{S}^2 + \lambda_{\text{BCS}}\hat{T}^+\hat{T}.$$

- I. Excitations are *similar* to the excitations in a disordered Fermi-gas.
- II. Small decay rate
- III. Substantial renormalizations



*Isn't it a Fermi liquid ?*

*Fermi liquid behavior follows from the fact that different wave functions are almost uncorrelated*


# Example 1: Coulomb Blockade




Coulomb Blockade



$N \rightarrow N+1$  transition





Coulomb Blockade Peak Spacing  
 Patel, et al. PRL 80 4522 (1998)  
 (Marcus Lab)


# Example 2: Spontaneous Magnetization

1. Disorder  
( $\times$ impurities)

2. Complex  
geometry


3.  $e$ - $e$  interactions

*chaotic  
one-particle  
motion*





## Example 2: Spontaneous Magnetization

1. Disorder (× impurities)
  2. Complex geometry
  3. e-e interactions
- chaotic one-particle motion*



Q • *What is the **spin** of the Quantum Dot in the ground state?*

# How to measure the Magnetization – **motion** of the Coulomb blockade peaks in the **parallel** magnetic field



In the presence of magnetic field

$$\hat{H}_{\text{int}} = \sum_{\alpha} \varepsilon_{\alpha} n_{\alpha} + J \hat{S}^2 + \vec{B} \cdot \hat{S}$$

Scaling:

the probability to find a ground state at a given magnetic field,  $B$ , with a given spin,  $S$ , depends on the combination rather than on  $B$  and  $J$  separately



$$X = J + g\mu_B \frac{B}{2S}$$



*Probability to observe a triplet state as a function of the parameter  $X$*

● - *results of the calculation based on the universal Hamiltonian with the RM one-particle states*

*The rest – exact diagonalization for Hubbard clusters with disorder. No adjustable parameters*



$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{int}}$$

$$\hat{H}_0 = \sum_{\alpha} \varepsilon_{\alpha} n_{\alpha}$$

$$\hat{H}_{\text{int}} = eV\hat{n} + E_c\hat{n}^2 + J\hat{S}^2 + \lambda_{\text{BCS}}\hat{T}^+\hat{T}.$$

- I. Excitations are *similar* to the excitations in a disordered Fermi-gas.
- II. *Small decay rate*
- III. *Substantial renormalizations*

## Small decay rate


- Why is it small
- What is it equal to
- What is the connection between the decay rate of the quasiparticles and the dephasing rate

Q ■ ■

?

# Quasiparticle decay rate at $T = 0$ in a *clean* Fermi Liquid.

I.  $d=3$



$$\frac{\hbar}{\tau_{e-e}(\epsilon)} \propto \left( \frac{\text{coupling}}{\text{constant}} \right)^2 \frac{\epsilon^2}{\epsilon_F} \quad d = 3$$

Reasons:


- At small  $\epsilon$  the energy transfer,  $\omega$ , is small and the integration over  $\epsilon'$  and  $\omega$  gives the factor  $\epsilon^2$ .
- The momentum transfer,  $q$ , is large and thus the scattering probability at given  $\epsilon'$  and  $\omega$  does not depend on  $\epsilon'$ ,  $\omega$  or  $\epsilon$

Quasiparticle decay rate at  $T = 0$  in a *clean* Fermi Liquid.

## II. Low dimensions

Small momenta transfer,  $q$ , become important at low dimensions because the scattering probability is proportional to the squared time of the interaction,

$$(\mathbf{q} \cdot \mathbf{v}_F)^{-2}$$



|   |   |         |
|---|---|---------|
|   | $\varepsilon^2 / \varepsilon_F$   | $d = 3$ |
| $\frac{\hbar}{\tau_{e-e}(\varepsilon)}$ | $\propto (\varepsilon^2 / \varepsilon_F) \log(\varepsilon_F / \varepsilon)$ | $d = 2$ |
|   | $\varepsilon$   | $d = 1$ |

Quasiparticle decay rate at  $T = 0$  in a *clean* Fermi Liquid.


### III. Applicability

|  |   |   |         |
|--|---|---|---------|
|  |   | $\varepsilon^2 / \varepsilon_F$   | $d = 3$ |
|  | $\frac{\hbar}{\tau_{e-e}(\varepsilon)}$ | $\propto (\varepsilon^2 / \varepsilon_F) \log(\varepsilon_F / \varepsilon)$ | $d = 2$ |
|  |   | $\varepsilon$   | $d = 1$ |

### Conclusions:

1. For  $d=3,2$  from  $\varepsilon \ll \varepsilon_F$  it follows that  $\varepsilon \tau_{e-e} \ll \hbar$ , i.e., that the **quasiparticles** are well determined and the Fermi-liquid approach is applicable.
2. For  $d=1$   $\varepsilon \tau_{e-e}$  is of the order of  $\hbar$ , i.e., that the Fermi-liquid approach is not valid for **1d** systems of interacting fermions.  
**Luttinger liquids**

Quasiparticle decay rate at  $T = 0$  in a *OD* Fermi Liquid.



Electronic spectrum is discrete

Need **offdiagonal** matrix elements

Quasiparticle decay is beyond the "universal Hamiltonian"

Quasiparticle decay rate is small as  $g^{-1}$

$$\tau_{ee}(\epsilon) \geq g \frac{\hbar}{\epsilon}$$

# CONCLUSIONS

One-particle chaos + moderate interaction of the electrons  $\mapsto$  to a rather simple Hamiltonian of the system, which can be called Zero-dimensional Fermi liquid.

The main parameter that justifies this description is the Thouless conductance, which is supposed to be large

Excitations are characterized by their one-particle energy, charge and spin, **but not by their momentum.**

These excitations have the lifetime, which is proportional to the Thouless conductance, i.e., is long.

This approach allows to describe Coulomb blockade (renormalization of the compressibility), as well as the substantial renormalization of the magnetic susceptibility and effects of superconducting pairing